

GENERALIZED LOW SOLUTION OF RT_k^1 PROBLEM

LU LIU

ABSTRACT. We study the "coding power" of an arbitrary RT_k^1 -instance. We prove that every RT_k^1 -instance admit non trivial generalized low solution. This is somewhat related to a problem proposed by Patey. We also answer a question proposed by Liu, i.e., we prove that there exists a $\mathbf{0}'$ -computable RT_3^1 -instance, I_3^1 , such that every RT_2^1 -instance admit a non trivial solution that does not compute any non trivial solution of I_3^1 .

1. INTRODUCTION

Reverse mathematics is a field that studies the proof theoretic strength of mathematical theorems. Many theorems are surprisingly found to be equivalent to one of the big five axioms [18]. Ramsey's theorem for pairs is probably one of the most famous exception.

Ramsey's theorem for single integers, RT_k^1 , is not interesting in the sense of reverse mathematics. Because over RCA_0 RT_k^1 is trivial. However, the "coding power" of an *arbitrary* instance of RT_k^1 attracts more and more attention since many admitting-homogenous-set theorems induced by binary relations reduce to the study of RT_k^1 instance. For example, RT_2^1, EM . Here EM is the Erdős-Moser theorem which says that every infinite tournament contains an infinite transitive subtournament (see also [8]).

In this paper, we prove two theorems about RT_k^1 instance. The first says that every instance of RT_k^1 admit generalized low solution. The second theorem prove the existence of a Muchnick degree of the solutions of an instance of RT_3^1 that can not be reduced to that of any instance of RT_2^1 . These results are of technical interest and are related to some recent arising questions in reverse mathematics as illustrated in subsection 1.1. We begin by introducing definitions of RT_k^n and review of the related literature.

Definition 1.1. Let $[X]^k$ denote $\{F \subseteq X : |F| = k\}$. A k -coloring on X is a function $c : [X]^k \rightarrow \{1, 2 \dots k\}$. A set $H \subseteq X$ is homogeneous for c iff c is constant on $[H]^k$. A stable 2-coloring is a function $c : [X]^2 \rightarrow \{1, 2\}$ such that there exists $i \in \{1, 2\}$ $|\{x \in X : c(x) \neq i\}| < \infty$.

Definition 1.2 (Ramsey's theorem [16]). (RT_k^n) For any n, k , every k -coloring of $[\mathbb{N}]^n$ admits an infinite homogeneous set.

The stable Ramsey's theorem for pair, SRT_k^2 , is RT_k^2 restricted to stable colorings.

Definition 1.3 (COH). Let $S_0, S_1, \dots$ be a sequence of sets. A set C is cohesive wrt $S_0, S_1, \dots$ iff $(\forall i \in \omega) |C \cap S_i| < \infty \vee |C \cap \bar{S}_i| < \infty$.

2010 *Mathematics Subject Classification.* Primary 68Q30 ; Secondary 03D32 03D80 28A78.

Key words and phrases. recursion theory, computability theory, reverse mathematics, RT_k^1 problem, Mathias forcing.

(COH): For every uniform sequence $S_0, S_1, \dots$ there exists an infinite cohesive set.

For more details see also [1] or [5]. There is a lot of literature on Ramsey theorems for pairs. To mention a few, Simpson [18] and Jockusch [7] proved that over RCA_0 , RT_k^n is equivalent to ACA_0 for $n > 2$. Jockusch [7] also showed that WKL_0 does not imply RT_2^2 . Seetapun and Slaman in their celebrated paper [17] proved that RT_2^2 does not imply ACA_0 over RCA_0 . Cholak, Jockusch and Slaman [1] proved that RT_2^2 is equivalent to $\text{SRT}_2^2 + \text{COH}$ over RCA_0 . Their paper also create one of the most important technique based on Mathias forcing, the *low*₂ construction. Liu [9],[10] separate RT_2^2 from WKL_0 and WWKL respectively. Therefore, combining with Jockusch [7] and the fact RT_2^2 is not equivalent to RCA_0 , RT_2^2 is not equivalent to any of the "big five". Most recently, Chong, Slaman and Yang [2] proved that SRT_2^2 does not imply COH and thus RT_2^2 . This settles a long standing problem. However, their model is nonstandard and thus leave the question that whether SRT_2^2 imply COH in standard arithmetic model. Another important progress is Patey and Yokoyama [15], they proved that over RCA_0 $\text{RT}_2^2 + \text{WKL}_0$ is $\tilde{\Pi}_3^0$ conservative. There they invent the notion of α -large etc that have prospective applications to other problems.

1.1. Main results. In this subsection we introduce our main results and the relation about our main results with the recent progress in reverse mathematics.

Patey [12] section 2 proposed several questions concerning computational complexity of solutions to Δ_2^0 instance of RT_2^1 . For example, "whether any Δ_2^0 RT_2^1 instance admit a solution that is both Δ_2^0 and *low*₂"; "whether there exists a Δ_2^0 RT_2^1 such that any Δ_2^0 non trivial solution of which is high". These questions are related to the currently most concerned problem in reverse mathematics, whether SRT_2^0 implies COH in ω -model. We here prove that any instance of RT_2^1 admit a generalized low solution.

Theorem 1.4. *For every instance of RT_2^1 , I_2^1 , there exists a non trivial solution to I_2^1 , G , such that $G' =_T G \oplus \mathbf{0}'$.*

Liu [10] proposed the question that whether there exists an instance of RT_3^1 such that of which the solution set is not Muchnik reducible to any solution set of any instance of RT_2^1 . We here give a positive answer in theorem 1.5. Similar results have been obtained independently by Dzhafarov [3], Hirschfeldt and Jockusch [6], Patey [14].

The question is of technical interest. Patey [13] proved that there exists a $\mathbf{0}'$ -computable instance of TT_2^1 , $I_{\text{TT}_2^1}$ such that the solution set of which is not Muchnik reducible to any instance of RT_2^1 . The proof relies on complexity of solution space of TT_2^1 , i.e., fix an arbitrary long initial segment of some instance of TT_2^1 , ρ , there exists sufficiently large M such that given any finitely many M -long initial segments of TT_2^1 -solutions of ρ , namely $\tau_i, i = 1, \dots, n$, there exists some extension of ρ , namely γ , such that any solution to any instance extending γ avoids $\tau_i, i \leq n$. However, this easy-avoidance property does not holds for RT_3^1 . RT_3^1 codes the solutions in a much more compact fashion. This is reflected by the fact that any instance of RT_3^1 computes a solution of itself. The proof of theorem 1.5 employs the method in Liu [10].

Theorem 1.5. *There exists a $\mathbf{0}'$ -computable instance of RT_3^1, I_3^1 , such that for every instance of RT_2^1, I_2^1 , there exists a non trivial solution to I_2^1, G_2^1 such that G_2^1 does not compute any non trivial solution to I_3^1 .*

Actually, by the proof of theorem 1.5, it is plain to see that,

Theorem 1.6. *There exists a $\mathbf{0}'$ -computable instance of RT_k^1, I_k^1 , such that for every instance of RT_{k-1}^1, I_{k-1}^1 , there exists a non trivial solution to I_{k-1}^1, G_{k-1}^1 such that G_{k-1}^1 does not compute any non trivial solution to I_k^1 . Where $k-1 \geq 1$ is arbitrary.*

The rest of this paper is devoted to the proof of the two theorem. We also propose new problems in section 5

2. NOTATIONS

We write $\rho * \tau$ to denote the string that concatenate τ after ρ . We sometimes regard a binary string $\rho \in 2^{<\omega}$ as a set $\{j : \rho(j) = 1\}$, and write $\rho \subseteq X$ for set containing relation, $\rho - \tau$ for set minus operation, $\rho \cap X$ for set intersection operation. For a sequence of string, $\dots \rho^i \subset \rho^{i+1} \dots, i \in \omega$, we write $\bigcup_{i \in \omega} \rho^i$ for the

string X such that $(\forall i \in \omega) X \upharpoonright_1^{\rho^i} = \rho^i$. When $\rho \cup \tau$ denote set union, we make assertion. We write $\rho \subset Y$ iff ρ is prefix of Y . *Empty string* is denoted by ε .

For a tree T , $[T]$ is the set of paths through T .

For a set X we write X' for the canonical jump, i.e., $X'(n) = 1$ iff $\Phi_n^X(n) \downarrow$. We write $\mathbf{0}'$ for the jump (Turing) degree.

3. PROOF OF THEOREM 1.4

3.1. Forcing conditions. Firstly, recall the Mathias forcing.

Definition 3.1.

- A Mathias condition is a pair (σ, X) with $\sigma \in 2^{<\omega}$ and $X \in 2^\omega$.
- (τ, Y) extends the Mathias condition (σ, X) iff $\sigma \subset \tau$ and $Y/\tau \subseteq X/\sigma$.
Write $(\tau, Y) \leq (\sigma, X)$ to denote the extension relation.
- A set G satisfies the Mathias condition (σ, X) if $\sigma \subset G$ and $G \subseteq X/\sigma$.

We say string $X \in 2^\omega$ codes an *ordered k -partition* iff

- $X = X_0 \oplus X_1 \oplus \dots \oplus X_{k-1}$ and
- $\bigcup_{i=0}^{k-1} X_i = \omega$

A class P is a k -partition class iff $\forall X \in P, X$ codes an ordered k -partition.

Definition 3.2. In the proof of theorem 1.4, a forcing condition is a tuple, $((\rho_{1,l}, \rho_{1,r}), \dots, (\rho_{k,l}, \rho_{k,r}), P, k)$, where $k > 0$ indicates the number of partitions and P is a Π_1^0 k -partition class.

Definition 3.3. We say set G *satisfies* condition $((\rho_{1,l}, \rho_{1,r}), \dots, (\rho_{k,l}, \rho_{k,r}), P, k)$ part j left side iff there exists $X_1 \oplus \dots \oplus X_k \in P$ such that G satisfies $(\rho_{i,l}, X_i)$. Similarly for right side.

We say set G *satisfies* condition c on part j iff it satisfies condition c part j on left or right side.

Definition 3.4. We say condition $d = ((\tau_{1,l}, \tau_{1,r}), \dots, (\tau_{m,l}, \tau_{m,r}), Q, m)$ *extends* condition

$c = ((\sigma_{1,l}, \sigma_{1,r}), \dots, (\sigma_{k,l}, \sigma_{k,r}), P, k)$, denoted by $d \leq c$, iff there is a function $f : m \rightarrow k$ satisfying $(\forall i \leq m \ \forall Y_1 \oplus \dots \oplus Y_m \in Q) (\exists X_1 \oplus \dots \oplus X_k \in P)$ such that $(\tau_{i,l}, Y_i) \leq (\sigma_{f(i),l}, X_{f(i)}) \wedge (\tau_{i,r}, Y_i) \leq (\sigma_{f(i),r}, X_{f(i)})$.

We say that

- f witnesses this extension;
- part i of d refines part $f(i)$ of c .

Definition 3.5. Part i of condition $((\rho_{1,l}, \rho_{1,r}), \dots, (\rho_{k,l}, \rho_{k,r}), P, k)$ is *acceptable* iff $(\exists X_1 \oplus \dots \oplus X_k \in P) |X_i| = \infty$

3.2. Outline. We will construct a sequence of forcing conditions $c_0 \geq c_1 \geq \dots \geq c_i \geq \dots$, together with a $\mathbf{0}'$ -computable function $F : (c_i, k) \mapsto (side, type)$. The function F tells how $c_{\langle e, r \rangle}$ satisfy requirement R_e, R_i , i.e., for every forcing condition $c_{\langle e, r \rangle}$ and part k of $c_{\langle e, r \rangle}$, $F(c_{\langle e, r \rangle}, k) = (left, 1)$ iff $\Phi_e^{\rho_{k,l}^{\langle e, r \rangle}}(e) \downarrow$; $F(c_{\langle e, r \rangle}, k) = (left, 0)$ iff for every G satisfying $c_{\langle e, r \rangle}$ on part k left side, $\Phi_e^G(e) \uparrow$ (lemma 3.7). Similarly for $F(c_{\langle e, r \rangle}, k) = (right, type)$.

The parts of these forcing conditions form a tree T . Nodes on level s of the tree represent the parts of condition c_s . Node j is a successor of node i iff for some s , j belongs to level $s+1$, i belongs to level s , and $f_{s+1}(j) = i$ where f_{s+1} is the witness of relation $c_{s+1} \leq c_s$ (see definition 3.4).

We will prove that for any instance Y , there exists a path along the forcing condition tree T , namely part r_i of condition c_i , $i \in \omega$, such that part r_i of c_i is acceptable and $\rho_{r_i,l}^i \subseteq A \wedge \rho_{r_i,r}^i \subseteq \bar{A}$ (subsection 3.4). And either $\bigcup_{j=1}^{\infty} \rho_{r_j,l}^j$ or

$\bigcup_{j=1}^{\infty} \rho_{r_j,r}^j$ is generalized low (subsection 3.5) and infinite (lemma 3.9).

To show the generalized low property, we prove that either for every e there exists r , $F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (left, \cdot)$; or for every r there exists e , $F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (right, \cdot)$ (lemma 3.10). Assume, without loss of generality, $\forall e \exists r F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (left, \cdot)$. Then we prove that given $G_l = \bigcup_{j=1}^{\infty} \rho_{r_j,l}^j$ we can $\mathbf{0}'$ -compute the path along T , i.e., the function $i \mapsto r_i$ (section 3.5). Given e , to decide whether $\Phi_e^{G_l}(e)$ halt, simply find r such that $F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (left, z)$. Finally, we prove that if $\forall e \exists r F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (left, \cdot)$, then $|G_l| = \infty$ (lemma 3.11).

3.3. Constructing forcing conditions and F . We start with condition $c_{-1} = ((\varepsilon, \varepsilon), \{\omega\}, 1)$. Given condition c_i , we show how to construct c_{i+1} . Suppose $i+1 = \langle e, r \rangle$. We will construct a sequence of forcing conditions $c_{i,1} \geq c_{i,2} \geq \dots \geq c_{i,k} = c_{i+1}$, where k is the number of parts of c_i , each dealing with a part of c_i to guarantee the successor of that part of c_{i+1} forces R_e, R_r as described in lemma 3.7.

Now we construct $c_{i,1}$. Consider the following Π_1^0 class,

$$(1) \quad [T^{c_i,1}] = \{X_{1l} \oplus X_{1r} \oplus X_2 \oplus \dots \oplus X_k : (X_{1l} \cup X_{1r}) \oplus X_2 \oplus \dots \oplus X_k \in P_i, \\ (\forall Z) \Phi_e^{(Z \cap X_{1l})/\rho_{1,l}^i}(e) \uparrow \wedge \Phi_r^{(Z \cap X_{1r})/\rho_{1,r}^i}(r) \uparrow\}$$

We divide into two cases, (1) $[T^{c_i,1}] = \emptyset$; (2) $[T^{c_i,1}] \neq \emptyset$.

If $[T^{c_i,1}] \neq \emptyset$ (in this case we adopt type 0 extension), split part 1 of c_i into two parts with identical initial segment $(\rho_{1,l}^i, \rho_{1,r}^i)$, replace P_i with $[T^{c_i,1}]$, i.e.,

$$c_{i,1} = ((\rho_{1,l}^i, \rho_{1,r}^i), (\rho_{1,l}^i, \rho_{1,r}^i), (\rho_{2,l}^i, \rho_{2,r}^i), (\rho_{3,l}^i, \rho_{3,r}^i), \dots, (\rho_{k,l}^i, \rho_{k,r}^i), [T^{c_i,1}], k+1)$$

. In this case, define $F(c_{i+1}, 1) = (\text{left}, 0)$, $F(c_{i+1}, 2) = (\text{right}, 0)$.

Else if $[T^{c_i,1}] = \emptyset$ (in this case we adopt type 1 extension), by compactness, there exists n such that for all $X = X_1 \oplus \dots \oplus X_k \in P_i$, $\rho \in \{0, 1\}^n$ there exists $\tau \in \{0, 1\}^n$ such that either $\tau \subseteq \rho \cap X_1 \wedge \Phi_e^{\rho_{1,l}^i * \tau}(e) \downarrow$ or $\tau \subseteq \bar{\rho} \cap X_1 \wedge \Phi_r^{\rho_{1,r}^i * \tau}(r) \downarrow$. Since $P_i \neq \emptyset$, fix an arbitrary $X \in P_i$. Suppose $\tau_1, \dots, \tau_{2^n}$ cover all $\rho \in \{0, 1\}^n$ in above way witnessed by $X = X_1 \oplus \dots \oplus X_k$. If $\tau_j \subseteq \rho \cap X_1 \wedge \Phi_e^{\rho_{1,l}^i * \tau}(e) \downarrow$ let $\rho_{j,l}^{i,1} = \rho_{1,l}^i * \tau_j$, $\rho_{j,r}^{i,1} = \rho_{1,r}^i * 0^n$ and define $F(c_{i+1}, j) = (\text{left}, 1)$; else if $\tau_j \subseteq \bar{\rho} \cap X_1 \wedge \Phi_r^{\rho_{1,r}^i * \tau}(r) \downarrow$ let $\rho_{j,l}^{i,1} = \rho_{1,l}^i * \tau_j$, $\rho_{j,r}^{i,1} = \rho_{1,r}^i * 0^n$ and define $F(c_{i+1}, j) = (\text{right}, 1)$. To construct $c_{i,1}$, split part 1 of c_i into 2^n many parts, and concatenate 0^n to initial segments of other parts of c_i , i.e., extend $\rho_{j,l}^i, \rho_{j,r}^i$ to $\rho_{j,l}^{i,1} * 0^n, \rho_{j,r}^{i,1} * 0^n$ for all $j \neq 1$. Furthermore, shrink P_i to $Z_1 \oplus \dots \oplus Z_k$ such that $Z_1 \supseteq \bigcup_{j=1}^{2^n} \tau_j$ and replicate part 1 of the shrunked P_i for 2^n many times, i.e.,

(2)

$$P_{i,1} = \left\{ \underbrace{Z_1 \oplus \dots \oplus Z_1}_{2^n \text{ many}} \oplus Z_2 \oplus Z_3 \oplus \dots \oplus Z_k : Z_1 \oplus Z_2 \oplus \dots \oplus Z_k \in P_i \wedge Z_1 \supseteq \bigcup_{j=1}^{2^n} \tau_j \right\}$$

(Here $\bigcup_{j=1}^{2^n} \tau_j$ is regarded as a set.) In summary,

$$c_{i,1} = ((\rho_{1,l}^{i,1}, \rho_{1,r}^{i,1}), \dots, (\rho_{2^n,l}^{i,1}, \rho_{2^n,r}^{i,1}), \dots, (\rho_{2^n+k-1,l}^{i,1}, \rho_{2^n+k-1,r}^{i,1}), P_{i,1}, 2^n + k - 1)$$

Where for all $j > 2^n$, $\rho_{j,l}^{i,1} = \rho_{j-2^n+1,l}^i * 0^n$, $\rho_{j,r}^{i,1} = \rho_{j-2^n+1,r}^i * 0^n$. It is clear that $P_{i,1} \neq \emptyset$ since it at least contains $X = X_1 \oplus \dots \oplus X_k$.

Remark 3.6. The purpose to concatenate 0^n to initial segments of the other parts is to enable $\mathbf{0}' \oplus G$ - compute the path along the forcing condition tree, i.e., the function $i \mapsto r_i$ (see lemma 3.9 and subsection 3.5). Note that in this way, all initial segments a forcing condition are of identical length.

It is clear that $c_{i,1} \leq c_i$. The forcing condition $c_{i,1}$ dealt with part 1 of c_i . After $c_{i,1}$ is constructed, construct $c_{i,2} \geq c_{i,3} \geq \dots \geq c_{i,k}$ similarly to deal with part 2, 3, $\dots$, k of c_i .

The following lemma 3.7 says that function F tells how forcing conditions $c_i, i \in \omega$, satisfy the requirements.

Lemma 3.7. *The function F is $\mathbf{0}'$ -computable and*

- If $F(c_{<e,r>}, j) = (\text{left}, 1)$, then for all G satisfying c_i on part j left side, i.e., $G \subseteq X_j / \rho_{j,l}^i \wedge G \supset \rho_{j,l}^i$, $\Phi_e^G(e) \downarrow$;
Similarly,
- If $F(c_{<e,r>}, j) = (\text{left}, 0)$, then for all G satisfying c_i on part j left side, $\Phi_e^G(e) \uparrow$;
- If $F(c_{<e,r>}, j) = (\text{right}, 1)$, then for all G satisfying c_i on part j right side, $\Phi_r^G(r) \downarrow$;

- If $F(c_{<e,r>,j}) = (\text{right}, 0)$, then for all G satisfying c_i on part j right side, $\Phi_r^G(r) \uparrow$;

Proof. The four items are obvious due to the construction of c_i . Note that the construction of c_i and F is uniform in $\mathbf{0}'$. Thus, $F \leq_T \mathbf{0}'$. $\square$

To construct the set G , we need the following lemma, which says that the forcing condition tree is built along all instances of RT_2^1 .

Lemma 3.8. *For any instance of RT_2^1 , Y , any forcing condition c_i there exists $j \in \omega$, parts $k_1, \dots, k_j$ of c_i with $(\forall s \leq j) \rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \overline{Y}$ such that*

$$(\forall X = X_1 \oplus \dots \oplus X_k \in P_i) \bigcup_{s=1}^j X_{k_s} = \omega$$

Proof. The proof is done by induction. Clearly, the lemma holds for c_{-1} . Assume it holds for c_i . We show that it holds for $c_{i,1}$. Fix an arbitrary instance of RT_2^1 , Y .

If $c_{i,1}$ is type 1 extension of c_i , as in section 3.3. Suppose for all $X = X_1 \oplus \dots \oplus X_k \in P_i$, $X_{k_1} \cup X_{k_2} \cup \dots \cup X_{k_j} = \omega$ and $\rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \overline{Y}$, with $(\forall s) k_s \neq 1$. Then it is obvious that for all $X = X_1 \oplus \dots \oplus X_{2^n} \oplus \dots \oplus X_{2^n+k-1} \in P_{i,1}$, $X_{2^n+k-1} \cup X_{2^n+k-2} \cup \dots \cup X_{2^n+k-j-1} = \omega$, $\rho_{2^n+k-1,l}^i \subseteq Y \wedge \rho_{2^n+k-1,r}^i \subseteq \overline{Y}$ for $s = 1, 2, \dots, j$ since there exists $Z_1 \oplus Z_2 \oplus \dots \oplus Z_k \in P_i$, $n \in \omega$ such that $Z_{k_s} = X_{2^n+k-1}$, and $\rho_{2^n+k-1,l}^i = \rho_{k_s,l}^i * 0^n \wedge \rho_{2^n+k-1,r}^i = \rho_{k_s,r}^i * 0^n$ for $s = 1, 2, \dots, j$.

Suppose for all $X = X_1 \oplus \dots \oplus X_k \in P_i$, $X_{k_1} \cup X_{k_2} \cup \dots \cup X_{k_j} = \omega$ and $(\forall s \leq j) \rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \overline{Y}$ with $(\forall s > 1) k_s \neq 1, k_1 = 1$. Take $\rho = Y \upharpoonright_{|\rho_{1,l}^i|+n} \in \{0,1\}^n$ and suppose τ_h covers ρ witnessed by X (recall the construction during type 1 extension), i.e., $\tau_h \subseteq \rho \cap X_1 \vee \tau_h \subseteq \overline{\rho} \cap X_1$. Then $\rho_{h,l}^{i,1} \subseteq Y \wedge \rho_{h,r}^{i,1} \subseteq \overline{Y}$ since $\tau_h \subseteq \rho \cap X_1 \rightarrow \rho_{h,l}^{i,1} = \rho_{1,l}^i * \tau_h \wedge \rho_{h,r}^{i,1} = \rho_{h,r}^i * 0^n$ and $\tau_h \subseteq \overline{\rho} \cap X_1 \rightarrow \rho_{h,r}^{i,1} = \rho_{1,r}^i * \tau_h \wedge \rho_{h,l}^{i,1} = \rho_{h,l}^i * 0^n$. But for any $X = X_1 \oplus \dots \oplus X_{2^n} \oplus \dots \oplus X_{2^n+k-1} \in P_{i,1}$ there exists $Z_1 \oplus Z_2 \oplus \dots \oplus Z_k \in P_i$ with $Z_1 = X_1 = X_2 = \dots = X_{2^n}$, $Z_{2^n+k-1} = X_{k_s}$ for $s = 2, 3, \dots, j$. Therefore $X_h \cup X_{2^n+k-1} \cup \dots \cup X_{2^n+k-j-1} = \omega$. And clearly $\rho_{h,l}^{i,1} \wedge \rho_{2^n+k-1,l}^{i,1} \subseteq Y \wedge \rho_{h,r}^{i,1} \wedge \rho_{2^n+k-1,r}^{i,1} \subseteq \overline{Y}$ for $s = 2, 3, \dots, k$.

If $c_{i,1}$ is type 0 extension of c_i . Suppose for all $X = X_1 \oplus \dots \oplus X_k \in P_i$, $X_{k_1} \cup X_{k_2} \cup \dots \cup X_{k_j} = \omega$ and $(\forall s \leq j) \rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \overline{Y}$, $(\forall s) k_s \neq 1$. Then it is obvious that for all $X = X_1 \oplus X_2 \oplus \dots \oplus X_{k+1} \in P_{i,1}$, $X_{k_1+1} \cup X_{k_2+1} \cup \dots \cup X_{k_j+1} = \omega$, $(\forall s) \rho_{k_s+1,l}^{i,1} \subseteq Y \wedge \rho_{k_s+1,r}^{i,1} \subseteq \overline{Y}$ since $Z_1 \cup Z_2 \oplus \dots \oplus Z_k \in P_i$ and $\rho_{k_s+1,l}^{i,1} = \rho_{k_s,l}^i \wedge \rho_{k_s+1,r}^{i,1} = \rho_{k_s,r}^i$ for $s = 1, 2, \dots, k$.

Suppose for all $X = X_1 \oplus \dots \oplus X_k \in P_i$, $X_{k_1} \cup X_{k_2} \cup \dots \cup X_{k_j} = \omega$ and $(\forall s \leq j) \rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \overline{Y}$, $(\forall s > 1) k_s \neq 1, k_1 = 1$. Let $Z_1 \oplus Z_2 \oplus \dots \oplus Z_{k+1} \in P_{i,1}$ be arbitrary. But clearly $\rho_{1,l}^{i,1} = \rho_{2,l}^{i,1} = \rho_{1,l}^i \subseteq Y \wedge \rho_{1,r}^{i,1} = \rho_{2,r}^{i,1} = \rho_{1,r}^i \subseteq \overline{Y}$, $\rho_{k_s+1,l}^{i,1} = \rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s+1,r}^{i,1} = \rho_{k_s,r}^i \subseteq \overline{Y}$ for $s = 2, 3, \dots, k_j$ and $Z_1 \cup Z_2 \cup Z_{k_2+1} \cup \dots \cup Z_{k_j+1} = \omega$, since $Z_1 \cup Z_2 \oplus Z_3 \oplus Z_4 \oplus \dots \oplus Z_{k+1} \in P_i$. $\square$

Later we need to prove that given $\bigcup_{i \in \omega} \rho_{r_i,l}^i$ or $\bigcup_{i \in \omega} \rho_{r_i,r}^i$ we can compute the path through forcing condition tree T , i.e., function $i \mapsto r_i$ (see subsection 3.5). This needs the following auxiliary lemma 3.9,

Lemma 3.9. *For any $i, I \in \omega$, any part k of c_i and any part K of c_I that is a decent of part k of c_i , if $\rho_{K,l}^I - \rho_{k,l}^i \neq \emptyset$, then for any part $k' \neq k$ of c_i and any part K' of c_I that is a decent of k' of c_i , we have, $\rho_{K',l}^I$ is incomparable with $\rho_{K,l}^I$.*

Proof. This is simply because the only chance for an initial segment to add new element is through type 1 extension. But whenever some initial segment is extended to $\rho * \tau$ with $\tau \neq \emptyset$ through type 1 extension during construction of c_s , $s > i$, each initial segment of the other parts of c_{s-1} , say ρ' , is extended to $\rho' * 0^{|\tau|}$. So $\rho * \tau$ is incomparable with any $\rho' * 0^{|\tau|}$, i.e., any initial segments whose parts in c_{s-1} is different with that of ρ . $\square$

3.4. Constructing G . Given instance A of RT_2^1 , to construct $G \subseteq A \vee G \subseteq \overline{A}$ that is generalized low, note that by lemma 3.8, each forcing condition c_i admit some part k that is acceptable and $\rho_{k,l}^i \subseteq A \wedge \rho_{k,r}^i \subseteq \overline{A}$. Also note that if part K of c_I is acceptable and part k of c_i is a parent node of part K of c_I , then part k of c_i is also acceptable. Therefore, the acceptable parts of all forcing conditions c_i , $i \in \omega$ form an infinite subtree of the whole forcing conditions tree T . Thus, the subtree admit a path, say part r_i of c_i , $i \in \omega$. Consider $G_l = \bigcup_{i \in \omega} \rho_{r_i,l}^i$, $G_r = \bigcup_{i \in \omega} \rho_{r_i,r}^i$. It is obvious that $G_l \subseteq A \wedge G_r \subseteq \overline{A}$. We will prove that either G_l or G_r is generalized low and infinite.

It is plain to see that,

Lemma 3.10. *Either $(\forall e)(\exists r)F(c_{<e,r>}, r_{<e,r>}) = (left, \cdot)$ or $(\forall r)(\exists e)F(c_{<e,r>}, r_{<e,r>}) = (right, \cdot)$.*

We use lemma 3.10 to prove that at least one of G_l, G_r is infinite.

Lemma 3.11. *Assume for all i part r_i of c_i is acceptable.*

We have, if $(\forall e)(\exists r)F(c_{<e,r>}, r_{<e,r>}) = (left, \cdot)$ then $|G_l| = \infty$.

Similarly, if $(\forall r)(\exists e)F(c_{<e,r>}, r_{<e,r>}) = (right, \cdot)$ then $|G_r| = \infty$.

Proof. Assume $(\forall e)(\exists r)F(c_{<e,r>}, r_{<e,r>}) = (left, \cdot)$. Consider such Turing functional E , $\Phi_E^Z(E) \downarrow$ if and only if $|Z| > E$. Let R be such that $F(c_{<E,R>}, r_{<E,R>}) = (left, z)$. Note that by the construction of $c_{<E,R>}$ and F , we have, either for all G satisfying $c_{<E,R>}$ part $r_{<E,R>}$ left side, $\Phi_E^G(E) \downarrow$; or for all G satisfying $c_{<E,R>}$ part $r_{<E,R>}$, left side $\Phi_E^G(E) \uparrow$, depending on $z = 1, 0$. But part $r_{<E,R>}$ of $c_{<E,R>}$ is acceptable. So there exists H satisfying $c_{<E,R>}$ on part $r_{<E,R>}$ left side that is infinite. Thus, $\Phi_E^H(E) \downarrow$ by definition of $\Phi_E(E)$. This implies that for all G satisfying $c_{<E,R>}$ part $r_{<E,R>}$ left side, $\Phi_E^G(E)$ halt. In particular $\Phi_E^{G_l}(E)$ halt. This implies $|G_l| > E$. The proof is accomplished by noting that E is arbitrary. $\square$

In the following proof of theorem 1.4 we assume, without loss of generality, $(\forall e)(\exists r)F(c_{<e,r>}, r_{<e,r>}) = (left, \cdot)$. Thus, by lemma 3.11 $|G_l| = \infty$.

3.5. Compute G' . To compute G'_l . We firstly show that we can compute the function $i \mapsto r_i$ using G_l and $\mathbf{0}'$.

Given $i = \langle e, r \rangle$ to compute $r_{<e,r>}$, firstly find (uniformly in i) a Turing functional Φ_E such that $E > |\rho_{1,l}^i|$, $\langle E, s \rangle > i$ for all $s \in \omega$ and $\Phi_E^G(E) \downarrow$ iff $|G| > E$. By our assumption, $(\forall e)(\exists r)F(c_{<e,r>}, r_{<e,r>}) = (left, \cdot)$, there exists R such that $F(c_{<E,R>}, r_{<E,R>}) = (left, z)$. Let R be an arbitrary such integer.

We show that on level $\langle E, R \rangle$ of the forcing condition tree T , there exists part k such that $\rho_{k,l}^{\langle E, R \rangle} \subset G_l \wedge \rho_{k,l}^{\langle E, R \rangle} - \rho_{f(k, \langle E, R \rangle, i), l}^i \neq \emptyset$, furthermore for any part k' of $c_{\langle E, R \rangle}$ if $\rho_{k',l}^{\langle E, R \rangle} \subset G_l \wedge \rho_{k',l}^{\langle E, R \rangle} - \rho_{f(k', \langle E, R \rangle, i), l}^i \neq \emptyset$ then part k' of $c_{\langle E, R \rangle}$ is a decent of part r_i of c_i where part $f(k', \langle E, R \rangle, i)$ of c_i is the accent of part k' of $c_{\langle E, R \rangle}$. Thus, to compute r_i , simply find (effectively in $G_l \oplus \mathbf{0}'$) a number R and part k of $c_{\langle E, R \rangle}$, $\rho_{k,l}^{\langle E, R \rangle} \subset G_l \wedge \rho_{k,l}^{\langle E, R \rangle} - \rho_{f(k, \langle E, R \rangle, i), l}^i \neq \emptyset$. Then part r_i is simply $f(k, \langle E, R \rangle, i)$.

To prove existence of k , we show that $r_{\langle E, R \rangle}$ is such a part. As in the proof of lemma 3.11, it must holds that $F(c_{\langle E, R \rangle}, r_{\langle E, R \rangle}) = (left, 1)$ since part $r_{\langle E, R \rangle}$ of $c_{\langle E, R \rangle}$ is acceptable. Therefore $|\{t : \rho_{r_{\langle E, R \rangle}, l}^{\langle E, R \rangle}(t) = 1\}| \geq E$. But since $E > |\rho_{1,l}^i|$ therefore $\rho_{r_{\langle E, R \rangle}, l}^{\langle E, R \rangle} - \rho_{f(r_{\langle E, R \rangle}, \langle E, R \rangle, i), l}^i \neq \emptyset$.

Now we show that for any part k' of $c_{\langle E, R \rangle}$ if $\rho_{k',l}^{\langle E, R \rangle} \subset G_l \wedge \rho_{k',l}^{\langle E, R \rangle} - \rho_{f(k', \langle E, R \rangle, i), l}^i \neq \emptyset$ then part k' of $c_{\langle E, R \rangle}$ is a decent of part r_i of c_i . Due to lemma 3.9, $\rho_{k',l}^{\langle E, R \rangle}$ is incomparable with any $\rho_{k,l}^{\langle E, R \rangle}$ when part k of $c_{\langle E, R \rangle}$ is not a decent of part $f(k', \langle E, R \rangle, i)$ of c_i . But $\rho_{k',l}^{\langle E, R \rangle} \subset G_l$ implies $\rho_{k',l}^{\langle E, R \rangle}$ equals to $\rho_{r_{\langle E, R \rangle}, l}^{\langle E, R \rangle}$ thus comparable to $\rho_{r_{\langle E, R \rangle}, l}^{\langle E, R \rangle}$. Therefore part $r_{\langle E, R \rangle}$ of $c_{\langle E, R \rangle}$ is a decent of part $f(k', \langle E, R \rangle, i)$. But on level i of the forcing condition tree T , part $r_{\langle E, R \rangle}$ has the unique accent node that is part r_i of c_i . Therefore $f(k', \langle E, R \rangle, i) = r_i$.

To compute G'_l . Given e , to decide whether $\Phi_e^{G_l}(e)$ halt, simply compute (in $G_l \oplus \mathbf{0}'$) r and $r_{\langle e, r \rangle}$ such that $F(c_{\langle e, r \rangle}, r_{\langle e, r \rangle}) = (left, z)$. Then $\Phi_e^{G_l}(e) \downarrow$ iff $z = 1$ and $\Phi_e^{G_l}(e) \uparrow$ iff $z = 0$.

4. PROOF OF THEOREM 1.5

4.1. Forcing condition. The forcing condition we use in the proof of theorem 1.5 is not Π_1^0 partition class. The Π_1^0 partition class is replaced by a single low partition, i.e.,

$$((\rho_{1,l}, \rho_{1,r}), (\rho_{2,l}, \rho_{2,r}), \dots, (\rho_{k,l}, \rho_{k,r}), X_1 \oplus X_2 \oplus \dots \oplus X_k, k)$$

where $\bigcup_{j \leq k} X_j = \omega$ and $X = X_1 \oplus \dots \oplus X_k$ is low.

Definitions in section 3.1 such as " $c \leq d$ ", " G satisfy c on part j left side", "acceptable" can clearly be inherited.

4.2. Outline. We will construct uniformly in $\mathbf{0}'$ a sequence of forcing conditions $\dots c_i \geq c_{i+1} \dots$ together with a sequence of RT_3^1 instance initial segment $\dots \beta^i \subset \beta^{i+1} \dots$ such that for any instance of RT_2^1 , A , there exists G encoded by the forcing condition that does not compute any solution to $A_3^1 = \bigcup_{j=1}^{\infty} \beta^j$.

Note that here by constructing a forcing condition $c = ((\rho_{1,l}, \rho_{1,r}), \dots, (\rho_{k,l}, \rho_{k,r}), X, k)$ we mean not only to demonstrate the existence of c but also compute the Turing functional t, t' such that $\Phi_t^{\emptyset'} = X$, $\Phi_{t'}^{\emptyset'} = X'$. The purpose is to guarantee $\mathbf{0}'$ -computability of the function $i \mapsto \beta^i$.

The requirement each forcing condition try to meet take the form as following.

4.1. $R_e(\beta) : (\forall n \leq |\beta|) \Phi_e^G(n) \downarrow \Rightarrow \Phi_e^G(n) = 0$ or Φ_e^G is trivial or Φ_e^G violate β deterministically (i.e., $\Phi_e^G(t) = \Phi_e^G(s) = 1 \wedge \beta(s) \neq \beta(t)$).

Note that if G satisfy all $R_e(\beta^e)$ for a sequence $\dots \beta^e \subset \beta^{e+1} \dots$, then G fail to compute any non trivial solution to $\bigcup_{j=1}^{\infty} \beta^j$.

We construct the forcing conditions and satisfy the requirements in the following way.

4.2. For any part k of $c_{\langle e, r \rangle}$, either for every G satisfying $c_{\langle e, r \rangle}$ on part k left side, G satisfies $R_e(\beta^{\langle e, r \rangle})$ or $R_e(\beta^{\langle e, r \rangle - 1})$; or for every G satisfying $c_{\langle e, r \rangle}$ on part k right side, G satisfies $R_r(\beta^{\langle e, r \rangle})$ or $R_r(\beta^{\langle e, r \rangle - 1})$.

Meanwhile, we guarantee a lemma 3.8 holds (see lemma 4.5).

Once such $c_i, \beta^i, i \in \omega$ are constructed, given an instance of RT_2^1 , A , there exists an infinite subtree of the forcing condition tree such that each node of the subtree represents an acceptable part and for every part k of c_i on that subtree, $\rho_{k,l}^i \subseteq A \wedge \rho_{k,r}^i \subseteq \bar{A}$. Thus, the infinite subtree admit a path, namely part r_i of c_i , $i \in \omega$ such that $\rho_{r_i,l}^i \subseteq A \wedge \rho_{r_i,r}^i \subseteq \bar{A}$. We show that either $G_l = \bigcup_{i \in \omega} \rho_{r_i,l}^i$ is infinite and satisfy all $R_e(\beta^e)$, $e \in \omega$; or $G_r = \bigcup_{i \in \omega} \rho_{r_i,r}^i$ is infinite and satisfy all $R_r(\beta^r)$, $r \in \omega$.

4.3. Constructing the forcing conditions and $\beta^i, i \in \omega$. We begin with some definitions which is also used in [10] [?]. We regard instances of RT_3^1 as functions $\omega \rightarrow \{1, 2, 3\}$.

Definition 4.3. For an instance of RT_3^1 , Y , we say Φ^X disagree with Y if and only if there exists $s, t \in \omega$ $\Phi^X(s) = \Phi^X(t) = 1$ and $Y(s) \neq Y(t)$.

As in the proof of theorem 1.4, let $c_{-1} = ((\varepsilon, \varepsilon), \{\omega\}, 1)$. $\beta^{-1} = \varepsilon$.

Suppose $i+1 = \langle e, r \rangle$. Given a condition $c_i = ((\rho_{1,l}^i, \rho_{1,r}^i), (\rho_{2,l}^i, \rho_{2,r}^i), \dots, (\rho_{k_i,l}^i, \rho_{k_i,r}^i), X_1^i \oplus \dots \oplus X_{k_i}^i, k_i)$, β^i , we construct a sequence of conditions $c_{i,1} \geq c_{i,2} \geq \dots \geq c_{i,k_i} = c_{i+1}$ together with $\beta^{i,1} \subset \beta^{i,2} \subset \dots \beta^{i,k_i} = \beta^{i+1}$, $c_{i,h}$ deals with part h of c_i to guarantee that part forces the requirement $R_e(\beta^{i,h})$ or $R_e(\beta^i)$, $R_r(\beta^{i,h})$ or $R_r(\beta^i)$ in the way mentioned in 4.2. In the following proof, we show how to deal with part 1 and construct $c_{i,1}$.

If $(\forall n \leq |\beta^i|) \Phi_e^{\rho_{1,l}^i}(n) \uparrow \vee \Phi_e^{\rho_{1,l}^i}(n) \downarrow = 0$ or $(\forall n \leq |\beta^i|) \Phi_r^{\rho_{1,r}^i}(n) \uparrow \vee \Phi_r^{\rho_{1,r}^i}(n) \downarrow = 0$, then we are done by letting $c^{i,1} = c^i$. Clearly $c^{i,1}$ part 1 forces $R_e(\beta^i)$, $R_r(\beta^i)$ as 4.2.

Assume in the following that there exists $n, m \leq |\beta^i|$ such that $\Phi_e^{\rho_{1,l}^i}(n) \downarrow = 1 \wedge \Phi_r^{\rho_{1,r}^i}(m) \downarrow = 1$. For an instance of RT_3^1 , Y , let

(3)

$$[T_Y^{c_{i,1}}] = \{X_{1l}^i \oplus X_{1r}^i \oplus X_2^i \oplus \dots \oplus X_{k_i}^i : X_{1l}^i \cup X_{1r}^i = X_1^i,$$

$$(\forall Z) \Phi_e^{(Z \cap X_{1l}^i) / \rho_{1,l}^i} \text{ does not disagree with } Y \wedge \Phi_r^{(Z \cap X_{1r}^i) / \rho_{1,r}^i} \text{ does not disagree with } Y\}$$

For an instance of RT_3^1 , Y , and $h \in \omega$ denote by $Y + h$ the function $\omega \rightarrow \{1, 2, 3\}$ $(Y + h)(n) = Y(n) + (h \bmod 3)$.

Consider the Π_1^{0, X^i} class

$$(4) \quad [T^{c_{i,1}}] = \{Y : [T_{Y/\beta^i}^{c_{i,1}}], [T_{Y+1/\beta^i}^{c_{i,1}}], [T_{Y+2/\beta^i}^{c_{i,1}}] \neq \emptyset\}$$

We divide into two cases (1) $[T^{c_i,1}] \neq \emptyset$; (2) $[T^{c_i,1}] = \emptyset$.

If $[T^{c_i,1}] \neq \emptyset$, then by low basis theorem, there exists X^i -low instance of RT_3^1 , Y such that $[T_{Y/\beta^i}^{c_i,1}], [T_{Y+1/\beta^i}^{c_i,1}], [T_{Y+2/\beta^i}^{c_i,1}] \neq \emptyset$. Note that $[T_Z^{c_i,1}]$ is a $\Pi_1^{0,X^i \oplus Z}$ class for any Z . Since Y is X^i -low so $Y+1, Y+2$ are also X^i -low. Therefore again, by low basis theorem, there exists a $X^i \oplus Y$ -low path through $T_{Y+h/\beta^i}^{c_i,1}$, for all $h = 0, 1, 2$, namely, $X_{1l}^{i,h} \oplus X_{1r}^{i,h} \oplus X_2^i \oplus \cdots \oplus X_{k_i}^i$.

To construct $c_{i,1}$ we apply *Cross* operation to $X_{1l}^{i,h} \oplus X_{1r}^{i,h} \oplus X_2^i \oplus \cdots \oplus X_{k_i}^i$, $h = 0, 1, 2$ (see also [10]), i.e.,

$$(5) \quad \begin{aligned} X^{i,1} = & (X_{1l}^{i,0} \cap X_{1l}^{i,1}) \oplus (X_{1l}^{i,1} \cap X_{1l}^{i,2}) \oplus (X_{1l}^{i,2} \cap X_{1l}^{i,0}) \\ & \oplus (X_{1r}^{i,0} \cap X_{1r}^{i,1}) \oplus (X_{1r}^{i,1} \cap X_{1r}^{i,2}) \oplus (X_{1r}^{i,2} \cap X_{1r}^{i,0}) \oplus X_2^i \oplus X_3^i \oplus \cdots \oplus X_{k_i}^i \end{aligned}$$

And replicate the initial segment $\rho_{1,l}^i, \rho_{1,r}^i$ for 6 times, i.e.,

$$c_{i,1} = (\underbrace{(\rho_{1,l}^i, \rho_{1,r}^i), \dots, (\rho_{1,l}^i, \rho_{1,r}^i), (\rho_{2,l}^i, \rho_{2,r}^i), (\rho_{3,l}^i, \rho_{3,r}^i), \dots, (\rho_{k_i,l}^i, \rho_{k_i,r}^i)}_{6 \text{ times}}, X^{i,1}, k_i+5)$$

Clearly $c_{i,1} \leq c_i$.

Note that,

- Since $X_{1l}^{i,h} \oplus X_{1r}^{i,h} \oplus X_2^i \oplus \cdots \oplus X_{k_i}^i$ is $X^i \oplus Y$ -low for $h = 0, 1, 2$ therefore $X^{i,1}$ is $X^i \oplus Y$ -low. And because Y is X^i -low, X^i is low, therefore $X^{i,1}$ is low. The construction is clearly uniform, thus we can $\mathbf{O}'$ compute (with input c_i, β^i) the Turing functional namely $t_{i,1}, t'_{i,1}$, such that $\Phi_{t_{i,1}}^{\mathbf{O}'} = X^{i,1}$, $\Phi_{t'_{i,1}}^{\mathbf{O}'} = (X^{i,1})'$.
- For every G satisfying $c^{i,1}$ on its first 3 parts left side, Φ_e^G is trivial; and for every G satisfying $c^{i,1}$ on its second 3 parts right side, Φ_r^G is trivial. This is because that if G satisfy the Mathias condition $(\rho_{1,l}^i, X_{1l}^{i,h} \cap X_{1l}^{i,g})$, then by definition of $[T_{Y+h/\beta^i}^{c_i,1}], [T_{Y+g/\beta^i}^{c_i,1}]$, Φ_e^G must be a solution of both $(Y+h)/\beta^i, (Y+g)/\beta^i$. However, $(Y+h)/\beta^i, (Y+g)/\beta^i$ share no common non trivial homogeneous set containing element in $\{1, 2, \dots, |\beta^i|\}$ while every G satisfying the Mathias condition $(\rho_{1,l}^i, X_{1l}^{i,h} \cap X_{1l}^{i,g})$, $(\exists n \leq |\beta^i|) \Phi_e^G(n) \downarrow = 1$.
- Because each element of X_1^i , m , there must exists $h \neq g \in \{0, 1, 2\}$ such that m is contained by either both $X_{1l}^{i,h}, X_{1l}^{i,g}$ or both $X_{1r}^{i,h}, X_{1r}^{i,g}$. Therefore,

$$(6) \quad \begin{aligned} & (X_{1l}^{i,0} \cap X_{1l}^{i,1}) \cup (X_{1l}^{i,1} \cap X_{1l}^{i,2}) \cup (X_{1l}^{i,2} \cap X_{1l}^{i,0}) \\ & \cup (X_{1r}^{i,0} \cap X_{1r}^{i,1}) \cup (X_{1r}^{i,1} \cap X_{1r}^{i,2}) \cup (X_{1r}^{i,2} \cap X_{1r}^{i,0}) = X_1^i \end{aligned}$$

If $[T^{c_i,1}] = \emptyset$, then there must exists some $Y \supset \beta^i$ such that $[T_Y^{c_i,1}] = \emptyset$. By compactness, there exists n , $Y \supset \beta^{i,1} \supset \beta^i$ for any $\rho \in \{0, 1\}^n$ there exists τ such that either $\tau \subseteq \rho \cap X_1 \wedge \Phi_e^{\rho_{1,l}^i * \tau}$ disagree with $\beta^{i,1}$ or $\tau \subseteq \bar{\rho} \cap X_1 \wedge \Phi_r^{\rho_{1,r}^i * \tau}$ disagree with $\beta^{i,1}$. Suppose $\tau_1, \dots, \tau_{2^n}$ cover all $\rho \in \{0, 1\}^n$ in above way. If $\tau_j \subseteq \rho \cap X_1 \wedge \Phi_e^{\rho_{1,l}^i * \tau}$ disagree with $\beta^{i,1}$, then let $\rho_{j,l}^{i,1} = \rho_{1,l}^i * \tau_j, \rho_{j,r}^{i,1} = \rho_{1,r}^i$; else if $\tau \subseteq \bar{\rho} \cap X_1 \wedge \Phi_r^{\rho_{1,r}^i * \tau}$ disagree with $\beta^{i,1}$, then let $\rho_{j,r}^{i,1} = \rho_{1,r}^i * \tau_j, \rho_{j,l}^{i,1} = \rho_{1,l}^i$. To construct $c_{i,1}$, split the initial segment of part 1 of c_i into 2^n many initial segments as above and preserve

all initial segments of other parts of c_i . Furthermore, replicate part 1 of X^i for 2^n many times, i.e.,

$$(7) \quad X^{i,1} = \underbrace{X_1 \oplus \cdots \oplus X_1}_{2^n \text{ many}} \oplus X_2 \oplus X_3 \oplus \cdots \oplus X_k$$

In summary,

$$c_{i,1} = ((\rho_{1,l}^{i,1}, \rho_{1,r}^{i,1}), \dots, (\rho_{2^n,l}^{i,1}, \rho_{2^n,r}^{i,1}), \dots, (\rho_{2^n+k-1,l}^{i,1}, \rho_{2^n+k-1,r}^{i,1}), X^{i,1}, 2^n + k - 1)$$

Where for all $j > 2^n$, $\rho_{j,l}^{i,1} = \rho_{j-2^n+1,l}^i$, $\rho_{j,r}^{i,1} = \rho_{j-2^n+1,r}^i$. It is clear that $c_{i,1} \leq c_i$ and $X^{i,1}$ is low.

The forcing condition $c_{i,1}$ dealt with part 1 of c_i . After $c_{i,1}$ is constructed, construct $c_{i,2} \geq c_{i,3} \geq \cdots \geq c_{i,k}$ similarly to deal with part 2, 3, $\dots$, k of c_i .

Similar to lemma 3.7, we can show that c_{i+1} satisfies the requirements $R_e(\beta^{i+1})$ or $R_e(\beta^i)$, $R_r(\beta^{i+1})$ or $R_r(\beta^i)$ as in 4.2.

Lemma 4.4. *For every e, r , for every part k of $c_{<e,r>}$,*

either for every G satisfying $c_{<e,r>}$ on part k left side, G satisfy $R_e(\beta^{<e,r>})$ or $R_e(\beta^{<e,r>-1})$;

or for every G satisfying $c_{<e,r>}$ on part k right side, G satisfy $R_r(\beta^{<e,r>})$ or $R_r(\beta^{<e,r>-1})$.

In the first case we say that c_{i+1} part k progresses on the left side and in the second case we say that c_{i+1} part k progresses on the right side.

To construct G , we establish the following lemma that is exactly the same as lemma 3.8.

Lemma 4.5. *For any instance of RT_2^1 , Y , any c_i , there exists $j \in \omega$, parts $k_1, \dots, k_j$ of c_i with $\rho_{k_s,l}^i \subseteq Y \wedge \rho_{k_s,r}^i \subseteq \bar{Y}$ such that*

$$\bigcup_{s=1}^j X_{k_s}^i = \omega$$

Proof. The proof concern 6 and proceeds exactly the same as 3.8. $\square$

Remark 4.6. In case $[T^{c_i,1}] = \emptyset$, differently with the proof of theorem 1.4, we need not concatenate 0^n to initial segments of other parts. Because given G we need not compute the path through the forcing condition tree, i.e., we do not need lemma 3.9 here. But we need to prove additionally that $A_3^1 = \bigcup_{j=1}^{\infty} \beta^j$ is $\mathbf{0}'$ -computable as in the following lemma 4.7.

Lemma 4.7. *The RT_3^1 instance $A_3^1 = \bigcup_{j=1}^{\infty} \beta^j$ is $\mathbf{0}'$ -computable.*

Proof. The proof is accomplished by noting that the construction is effective in $\mathbf{0}'$. $\square$

4.4. Constructing G . Let $A_3^1 = \bigcup_{j=1}^{\infty} \beta^j$. Given instance A of RT_2^1 , to construct

$G \subseteq A \vee G \subseteq \bar{A}$ that does not compute any non trivial solution to A_3^1 , note that by lemma 3.8, each forcing condition c_i admit some part k that is acceptable and $\rho_{k,l}^i \subseteq A \wedge \rho_{k,r}^i \subseteq \bar{A}$. Also note that if part K of c_I is acceptable and part k of c_i is

an accent node of part K of c_I , then part k of c_i is also acceptable. Therefore, the acceptable parts of all forcing conditions c_i , $i \in \omega$ form an infinite subtree of the whole forcing conditions tree T . Thus, the subtree admit a path, say part r_i of c_i , $i \in \omega$. Consider $G_l = \bigcup_{i \in \omega} \rho_{r_i, l}^i$, $G_r = \bigcup_{i \in \omega} \rho_{r_i, r}^i$. It is obvious that $G_l \subseteq A \wedge G_r \subseteq \bar{A}$.

We will prove that either G_l or G_r fails to compute any non trivial solution of A_3^1 and is infinite.

It is plain to see that,

Lemma 4.8. *Either $(\forall e)(\exists r) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on left side or $(\forall r)(\exists e) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on right side.*

(Recall the definition of progress on the paragraph after lemma 4.4)

We use lemma 4.8 to prove that at least one of G_l, G_r is infinite.

Lemma 4.9. *Assume for all i part r_i of c_i is acceptable.*

We have, if $(\forall e)(\exists r) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on left side, then $|G_l| = \infty$.

Similarly, if $(\forall r)(\exists e) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on right side, then $|G_r| = \infty$.

Proof. The proof goes almost the same as 3.11. Without loss of generality suppose $(\forall e)(\exists r) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on left side. For any c it suffices to prove that $|G_l| > c$. Consider such Turing functional e that for any X $\Phi_e^X(1) = 1$ and $(\forall n \neq 1) \Phi_e^X(n) \downarrow = 1$ iff $|X| > c$. Since there exists r such that $c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progress on the left while $\Phi_e^X(1) = 1$, therefore it must holds that either for any G satisfying $c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ left side Φ_e^G violate $\beta^{\langle e, r \rangle}$ explicitly or Φ_e^G is trivial. But $c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ is acceptable thus by definition of Φ_e , there exists G satisfying $c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ left side such that Φ_e^G is non trivial. Therefore Φ_e^G violate $\beta^{\langle e, r \rangle}$ explicitly for all G satisfying $c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ left side. This implies that there exists $n \neq 1$ such that $\Phi_e^{\rho_{r_{\langle e, r \rangle}, l}^i}(n) = 1$ which implies $|\rho_{r_{\langle e, r \rangle}, l}^i| > c$. $\square$

Lemma 4.8, 4.9 together proved that either $G_l \subseteq A$ is infinite and does not compute any non trivial solution of A_3^1 or $G_r \subseteq \bar{A}$ is infinite and does not compute any non trivial solution of A_3^1 . To see this, suppose $(\forall e)(\exists r) \ c_{\langle e, r \rangle}$ part $r_{\langle e, r \rangle}$ progresses on left side, we show that for any e $\Phi_e^{G_l}$ violate A_3^1 explicitly or is trivial. Given e there exists infinitely many Turing functional $\Phi_{e_j}, j \in \omega$ such that all $\Phi_{e_j}, j \in \omega$ are exactly the same as Φ_e . However, for every e_j there exists r_j such that G_l satisfy $R_{e_j}(\beta^{\langle e_j, r_j \rangle})$ or $R_{e_j}(\beta^{\langle e_j, r_j \rangle - 1})$. Therefore by definition of requirement 4.1 (also note that $\bigcup_{j \in \omega} \beta^{\langle e_j, r_j \rangle - 1} = A_3^1$) either $(\forall n \in \omega) \Phi_e^{G_l}(n) \downarrow \Rightarrow \Phi_e^{G_l}(n) = 0$ or $\Phi_e^{G_l}$ violate some $\beta^{\langle e_j, r_j \rangle}$ explicitly or $\Phi_e^{G_l}$ is trivial.

Thus the proof of theorem 1.5 is accomplished.

5. FURTHER DISCUSSION AND QUESTIONS

The results are of technical interest. The proof is different with that of [10] or [4] in the sense that here we construct forcing conditions along all instances of the problem (RT_2^1) . Where in [10] we construct the objective set along a single instance. In another words, we pre choose a path through the forcing condition tree during the construction, and need not look at the construction else where in [10]. But here, to construct a set of "homogeneous" solutions intersecting with all instances (of RT_2^1), it is necessary to construct the forcing conditions along all

instances simultaneously. The difference is reflected by the type 0 extension and lemma 3.9, 4.5.

The results in this paper and many results arising recently [8] [11] [19], which says there exists somewhat weak solution to some instance, motivate us to find pure combinatorial conditions for a problem to admit "weak" solution in all of its instance. For example, taking "weak" to be generalized low.

The proof that RT_2^1 admit generalized low solution in every instance is somewhat robust. Therefore, we wonder if many other problems also has this property.

Question 5.1. *Is there a purely combinatorial condition that is necessary and sufficient for a problem P to admit generalized low solution in all its instances?*

For many known problems, say (S)CAC, (S)ADS, (S) RT_2^2 etc, they either encode large functions (every function g there exists instance of the problem such that all non trivial solutions compute some $f \geq g$) thus encode hyperarithmetic degree or admit generalized low solution. It'd be interesting to see some counterexamples.

Question 5.2. *Does there exists a "natural" problem that neither encode large function nor admit generalized low solutions?*

REFERENCES

1. P.A. Cholak, C.G. Jockusch, and T.A. Slaman, *On the strength of Ramsey's theorem for pairs*, Journal of Symbolic Logic **66** (2001), no. 1, 1–55.
2. Chitai Chong, Theodore Slaman, and Yue Yang, *The metamathematics of stable ramseys theorem for pairs*, Journal of the American Mathematical Society **27** (2014), no. 3, 863–892.
3. Damir Dzhafarov, *Cohesive avoidance and strong reductions*, Proceedings of the American Mathematical Society **143** (2015), no. 2, 869–876.
4. Damir D. Dzhafarov and Carl G. Jockusch, *Ramsey's theorem and cone avoidance*, Journal of Symbolic Logic **74** (2009), 557–578.
5. Denis R Hirschfeldt, *Slicing the truth: On the computable and reverse mathematics of combinatorial principles*, World Scientific, 2014.
6. Denis R Hirschfeldt and Carl G Jockusch Jr, *On notions of computability theoretic reduction between π_1^1 Σ_1^1 principles*, To appear (2015).
7. Carl G Jockusch, *Ramsey's theorem and recursion theory*, The Journal of Symbolic Logic **37** (1972), no. 02, 268–280.
8. Manuel Lerman, Reed Solomon, and Henry Towsner, *Separating principles below ramsey's theorem for pairs*, Journal of Mathematical Logic **13** (2013), no. 02, 1350007.
9. Jiayi Liu, *RT_2^2 does not imply WKL_0* , Journal of Symbolic Logic **77** (2012), no. 2, 609–620.
10. Lu Liu, *Cone avoiding closed sets*, Transactions of the American Mathematical Society **367** (2015), no. 3, 1609–1630.
11. Ludovic Patey, *Iterative forcing and hyperimmunity in reverse mathematics*, arXiv preprint arXiv:1501.07709 (2015).
12. ———, *Open questions about ramsey-type statements in reverse mathematics*, arXiv preprint arXiv:1506.04780 (2015).
13. ———, *The strength of the tree theorem for pairs in reverse mathematics*, arXiv preprint arXiv:1505.01057 (2015).
14. ———, *The weakness of being cohesive, thin or free in reverse mathematics*, arXiv preprint arXiv:1502.03709 (2015).
15. Ludovic Patey and Keita Yokoyama, *The proof-theoretic strength of ramsey's theorem for pairs and two colors*, arXiv preprint arXiv:1601.00050 (2016).
16. F.P. Ramsey, *On a problem of formal logic*, Proceedings of the London Mathematical Society **2** (1930), no. 1, 264.
17. David Seetapun, Theodore A Slaman, et al., *On the strength of ramsey's theorem*, Notre Dame Journal of Formal Logic **36** (1995), no. 4, 570–582.
18. S.G. Simpson, *Subsystems of second order arithmetic*, Springer, 1998.
19. Wei Wang, *The definability strength of combinatorial principles, 2014. to appear.*

E-mail address: `g.jiayi.liu@gmail.com`

DEPARTMENT OF MATHEMATICS AND STATISTICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA,
410083, CHINA